JC20 Rec'd PCT/PTO 0-1 APR 2002

FORM PTO-1390

U.S. Department of Commerce Patent and Trademark Office

TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)

Attorney's Docket No. 2577-138

U.S. Application No. (if known one 17 CER 1 S)

CONCERNING A FILING UNDER 35 U.S.C. 371 10/089543 INTERNATIONAL APPLICATION NO. INTERNATIONAL FILING DATE PRIORITY DATE CLAIMED PCT/SG00/00111 August 1, 2000

TITLE OF INVENTION

ISOLATION AND CHARACTERIZATION OF A FIBER-SPECIFIC β-TUBULIN PROMOTER FROM COTTON

Lin CAI, Xuebao LI, Ninghui CHENG and Jian-Wei LIU

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

- [X] This is a FIRST submission of items concerning a filing under 35 U.S.C. 371
- 1 This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
- [] This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
- The US has been elected by the expiration of 19 months from the priority date (Article 31).
- f X 1 A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. [X] is attached hereto (required only if not communicated by the International Bureau).
 - b. [X] has been communicated by the International Bureau.
 - c. [] is not required, as the application was filed in the United States Receiving Office (RO/US)
 - An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. [] is attached hereto.
 - b. [] has been previously submitted under 35 U.S.C. 154(d)(4).
- 7. [X] Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. [] are attached hereto (required only if not communicated by the International Bureau). b. [] have been communicated by the International Bureau.
 - have not been made; however, the time limit for making such amendments has NOT expired.
 - [X] have not been made and will not be made.
- An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
- [] An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
- An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

ITEMS 11. TO 20. below concern other document(s) or information included:

- An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
- 12. [An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. A FIRST preliminary amendment.
 - A SECOND or SUBSEQUENT preliminary amendment.
 - 15. F A substitute specification.
 - 16 f A change of power of attorney and/or address letter.
 - A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825
 - 1 A copy of the published international application.
 - 19. [A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
- 20. [X] Other items or information: 3 Drawing Sheets; Application Data Sheet; Sequence Listing (paper copy): International Search Report

U.S. APPLICATION NO/UT B 9549 3 INTERNATIONAL APPLICATION NO. PCT/SG00/00111			ATTORNEY DOCKET NO. 2577-138		
21 [X The following fees are submitted: Basic National Fee (37 CFR 1.492)(a)(1)-(5): Rother international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and international Search Report Not Prepared by EPO or JPO. \$ 1,040.00 International Search Report has been prepared by the EPO or JPO. \$ 8890.00 International Search Report has been prepared by the EPO or JPO. \$ 8890.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search (67 GFR 1.445(a)(2)) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Saminary Samination fee (37 CFR 1.482) paid to USPTO Saminary Saminary				CALCULATIONS	PTQ USE ONLY
		TER APPROPRIATE BAS		\$ 890.00	
Surcharge of \$130.00 for fur months from the earliest clair	nishing the oath or d med priority date (37	eclaration later than [] CFR 1.492(e)).	20 [] 30	s	
Claims	Number Filed	Number Extra	Rate		
Total Claims	3 -20 =		X \$18.00	\$	
Independent Claims	3 - 3 ==		X \$84.00	s	
Multiple dependent claim(s)	(if applicable)		+ \$280.00	\$	
		TOTAL OF ABOVE CA	ALCULATIONS =	\$ 890.00	
Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
SUBTOTAL =				\$ 890.00	
Processing fee of \$130.00 for months from the earliest claim	Processing fee of \$130.00 for furnishing the English translation later than []20 []30 months from the earliest claimed priority date (37 CFR 1.492(f)).				
TOTAL NATIONAL FEE =				\$ 890.00	
Fee for recording the enclosed accompanied by an appropria	d assignment (37 CF te cover sheet (37 CI	R 1.21(h)). The assignment of the second of the R 3.28, 3.31). \$40.00 per second of the second of th	ent must be er property +	s	
		TOTAL FE	ES ENCLOSED =	\$ 890.00	
				Amount to be refunded	\$
				charged	\$
a. A check in the amount of \$890.00 to cover the above fees is enclosed. Please charge my Deposit Account No. 02-2135 in the amount of \$					
Washington, D.C. 20005 40,315 Phone: 2027/83-6040 Registration Number					

2577-138A MIB:clw

10

15

20

25

3/17/

Technical Field

The present invention relates to the field of plant molecular biology, in particular to transgenic plants and promoters useful in creating transgenic plants, and more particularly to fiber-specific promoters.

Background of the Invention

Cotton is the most extensively used natural fiber in the textile industry. Annual production of cotton worldwide is over 100 million bales valued at 45 billion U.S. dollars. Although significant improvements have been made in quality and yield of the fibers by means of classical breeding in the past decades, the potential for further improving fiber properties through classical breeding is limited due to requirements for species compatibility and available traits. Genetic engineering provides novel approaches for further improving cotton by introducing genes to create new germplasms with highly desirable characteristics.

Cotton fibers (seed hairs) are single-cell trichomes that undergo rapid and synchronous elongation. Cortical microtubules provide spatial information necessary for the alignment of cellulose microfibrils that confine and regulate cell elongation [Giddings and Staehelin, 1991; Cyr and Palevitz, 1995; Fisher and Cyr, 1995]. Fiber development consists of four overlapping stages (i.e. initiation, primary cell wall formation, secondary

cell wall formation and maturation) [Basra and Malik, 1984]. Tubulins and actins may play functionally important roles in developing fiber cells. Mature fiber is a biological composite of cellulose, water, small quantities of proteins, pectins, hemicellulose, mineral substances, wax, small amounts of organic 5 acids, sugars, and pigments that provides excellent wearability and aesthetics (Arthur, 1990; Basra and Malik, 1984; Ryser, 1985]. Many genes are required for the fiber differentiation and development. These genes are differentially expressed during 10 different stages of the fiber development, and so far only a few of the genes involved in the biosynthesis of the large numbers of fiber-specific structural proteins, enzymes, polysaccharides, waxes or lignins have been identified [John and Crow, 1992; John, 1996a; Song and Allen, 1997; Ma et al., 1997; Kawai et 15 al., 1998; Whittaker and Triplett, 1999]. These isolated genes may be considered as having potential application in cotton

fiber improvement due to the character of their fiber-specific expression. For example, John has been using fiber-specific gene promoters to produce genetically engineering cotton for

altered fibers [John, 1996b, 1997a, 1997b].

20

25

30

A promoter is a DNA fragment that determines temporal and spatial specificity of gene expression during plant and animal development. Many tissue-specific genes and their promoters were identified and isolated from a wide variety of plants and animals over the past decade, including some cotton tissue-specific genes and promoters (Loguerico et al., 1999; Kawai et al., 1998; Song and Allen, 1997; Ma et al, 1997; John, 1996a; Rinehart et al., 1996; Hasenfratz et al, 1995; John and Peterson, 1994; John and Crow, 1992). A few promoters have been shown to control gene expression in a fiber-specific manner in cotton (Rinehart et al., 1996; John, 1996a; John and Crow, 1992). Some plant tissue-specific promoters can be utilized to

express foreign proteins in specific tissues in a developmentally regulated pattern [John, 1996b, 1997a, 1997b]. Summary of the Invention

A fiber-specific gene (named CFTUB2), encoding β-tubulin, was isolated from cotton. The isolated complete CFTUB2 cDNA is 1.623 kb in length including 1.338 kb of open reading frame. Based on the CFTUB2 cDNA sequence, two CFTUB2 promoter fragments (1.433 kb and 0.984 kb) were isolated from cotton. The two CFTUB2 promoter fragments (1.3 and 0.9 kb) were fused with the GUS gene to construct gene expression vectors for analyzing the function of the promoter. Transgenic cotton and tobacco plants with the CFTUB2 promoter/GUS fusion genes were identified by Southern blot hybridization. In all the transgenic cotton plants studied, GUS activity was detected only in young fibers, but not in the flower organs such as anthers, petals and sepals, or in leaves and roots. This result, together with Northern blot analysis, indicates that the CFTUB2 promoter is fiber-specific in cotton. The promoter controls specific gene expression at the transcriptional level in cotton fibers. The isolated promoter may be used in improving cotton fibers to create new cotton varieties with high fiber quality and yield by gene manipulation.

Brief Description of the Figures

5

10

1.5

20

25

30

Figure 1 shows the nucleotide sequence of the cotton CFTUB2 gene cDNA (1623 bp; SEO ID NO: 1).

Figure 2 shows the nucleotide sequence of the isolated 1433 bp CFTUB2 promoter fragment (SEQ ID NO: 2. The 984 bp fragment corresponds to nucleotides 449-1433 of this sequence.

Figure 3 shows constructs of the CFTUB2 promoter fused with the gus gene in expression vectors.

Detailed Description

5

10

15

20

25

30

The CFTUB2 promoter is an active fiber-specific promoter in cotton. Results of a Northern blot analysis of cDNAs from a variety of cotton tissues showed that a cDNA clone comprising the CFTUB2 gene was strongly expressed in young fibers of 8 and 14 days postanthesis (DPA), and also expressed in young ovules of 4, 8 and 14 DPA, but less or not at all in other tissues. Sequencing of the cDNA clone revealed that it was 1623 bp in length containing a open reading frame of 1338 bp (Figure 1). Comparing the nucleotide and predicted polypeptide sequences of the cotton CFTUB2 with the data banks, it was found that the CFTUB2 cDNA shared 96%-98% homology at the amino acid level and over 78% homology at the nucleotide level with the known B-tubulin cDNAs and genes from other plants (such as Arabidopsis, tobacco, rice, soybean, maize, potato, carrot, etc.) [Liaud et al., 1992; Snustad et al., 1992; Villemur et al, 1994; Tonoike, et al., 1994; Taylor et al., 1994; Kang et al., 1994; Okamura et al., 1997; Chu et al., 1998; Okamura et al., 19991.

The transcripts of the CFTUB2 gene exhibited the highest accumulation in cotton young fibers of 8 DPA, and then there was a visible decrease in the accumulation of the gene products (mRNA) with further development of the fibers. Comparison of gene expression in different developmental stages of cotton ovules also showed that the gene transcripts accumulated more in 8 DPA ovules than in 4 and 14 DPA, and there was a gradual and visible decrease to an undetectable level in the accumulation of gene products with fiber development from 8 DPA to 28 DPA. This suggests that the gene is specifically expressed with a strict regulation at the transcriptional level during cotton fiber and ovule development, as with other cotton fiber-specific genes [Whittaker and Triplett, 1999; Shin and Brown, 1999; Kawai et

al., 1998; John, 1996a; Song and Allen, 1997; Ma et al, 1997; Rinehart et al., 1996; John and Crow, 1992].

5

10

15

20

25

30

Two fragments in the promoter region were isolated and cloned into pGEM-T vector, respectively. One fragment of the CFTUB2 promoter was 1433 bp in length (Figure 2), and another was 984 bp long. Both fragments functioned as active, fiberspecific promoters. The constructs of CFTUB2 promoter/GUS fusion gene were used to transform tobacco and cotton by Agrobacterium-mediated gene transfer, using the pBI121 vector containing CaMV35S promoter/GUS fusion as a positive control. Consistent with the results from Northern blot analysis, the GUS gene driven by CFTUB2 promoter specifically expressed in the young fibers, but not in other tissues, in all the 31 transgenic cotton plants studied, while the GUS activity was detected in all the tissues of positive control cotton plants (35S:GUS). A total of 36 transformed cotton plants were obtained and transplanted in soil to grow to maturation. Similarly, it was found that under the CFTUB2 promoter, GUS gene activity was only detected in the seeds in all of the 15 transgenic tobacco plants studied, suggesting the CFTUB2 promoter activity was also tissue-specific in tobacco (the cotton fiber, being an elongated hair of the seed coat, finds histological correspondence in the tobacco seed coat). This result, together with the above Northern blot analysis, indicates that the CFTUB2 promoter controls gene specific expression at the transcriptional level in cotton fibers.

Accordingly, one embodiment of the present invention is a fiber-specific promoter obtained from the cotton fiber $\beta\text{-tubulin}$ gene CFTUB2.

Another embodiment of the present invention is a fiberspecific promoter comprising a 1433 kb active fragment of the cotton fiber CFTUB2 gene promoter. Another embodiment of the present invention is a fiberspecific promoter comprising a 984 kb active fragment of the cotton fiber CFTUB2 gene promoter.

5

10

1.5

20

25

30

Stll another embodiment of the present invention is a promoter that is cotton fiber-specific comprising an active fragment of the CFTUB2 promoter fragment of SEQ ID NO: 2. An active fragment is a sequence of shorter length than SEQ ID NO: 2 which still retains activity as a fiber-specific promoter in cotton. A fragment can comprise excisions, deletions, truncations or substitutions of the sequence of SEQ ID NO: 2, or a combination of these. A preferred active fragment is the fragment consisting of nucleotides 449-1433 of SEO ID NO: 2.

The promoters of the present invention are useful in creating transgenic cotton having altered fiber characteristics. The use of the fiber-specific promoters of the present invention permits selective expression of a transgene in the cotton fiber, permitting greater latitude in the types of transgenes employed. Selective expression avoids problems such as the metabolic burden imposed on a transgenic plant by systemic expression of a transgene, or the adverse effects of the expression of a transgene in non-fiber tissues. Examples for expressing desirable genes in cotton fiber, but not in other parts of the cotton plants include: (1) anthocynin genes for colored cotton, (2) silk protein genes from silk worm or spiders for increased strength of cotton fiber, (3) and biosynthesis of polyhydroxybutrate in cotton fiber for improved thermal properties and insulating characteristics [John, et al., 1996]. There are numerous examples in the art of fiber-enhancing genes that could be advantageously linked to the promoters of the present invention, and used to transform cotton using well-known techniques (see, e.g., Umbeck, 1992), to achieve expression of

the transgene in transgenic cotton fibers. <u>See e.g.</u>, John, 1996b, 1997a, 1997b; John et al., 1996.

EXAMPLE 1: Isolation of fiber-specific cDNA encoding CFTUB2 sequences expressed early during fiber development of cotton

5

10

15

20

25

30

Cotton seeds were surface-sterilized with 70% ethanol for 30-60 seconds and 10% H_2O_2 for 30-60 minutes, followed by washing with sterile water. The seeds germinated on $\frac{1}{2}$ MS medium on light at 28°C in a culture room, and cotyledons and hypocotyls cut from sterile seedlings were used as transformation explant materials. Cotton plants were grown in pots for DNA and RNA extraction.

Total RNA was extracted from young fibers, ovaries, anthers, petals, sepals, leaves and roots of cotton by using the guanidinium thiocyanate method or SV Total RNA Isolation System (Promega). Poly(A)+RNA was purified by using oligo(dT)-cellulose spin columns from an mRNA purification kit (Pharmacia Biotech). Cotton cDNA was synthesized by using a cDNA synthesis kit (Pharmacia Biotech). Cotton cDNA libraries were constructed by inserting the cDNA fragments into the ZAP express vector (Stratagene).

Poly(A)+ RNAs from cotton young fibers of about 8 and 14 days postanthesis (DPA), respectively, were converted to cDNAs which were used to construct cotton cDNA libraries. From the fiber cDNA libraries, about 200 cDNA clones were randomly picked out and subsequently sequenced. Some clones with potential involvement in cell expansion were selected according to the sequence data.

To find cDNA clones whose transcripts are specifically expressed in cotton fibers, the expression pattern of the $\,$

selected cDNA clones was analyzed by Northern blot hybridization with total RNAs isolated from cotton fibers, ovules, anthers, petals, sepals, squares, leaves and roots, using probes from the clones. RNA samples from the different cotton tissues were separated on agarose-formaldehyde gels, and transferred onto Hybond-N nylon membranes by capillary blotting. RNA Northern blots were hybridized in ExpressHyb solution (Clontech) at 68°C with ³²P cDNA probes prepared by random labeling (Promega Prime-a-Gene Labeling System). After hybridization, the blots were washed at 68°C in 0.1 x SSC, 0.5% SDS for 30-60 minutes. The experimental results showed that one cDNA clone strongly expressed in young fibers of 8 and 14 DPA, and also expressed in young ovules of 4, 8 and 14 DPA, but less or not at all in other tissues.

5

10

15

20

25

3.0

PCR fragments and cDNA fragments were subcloned into vectors, and plasmid DNA prepared with a Qiagen plasmid kit was used as templates in PCR reactions. The PCR products were sequenced by autosequencer. Sequencing of the cDNA clone revealed that it was 1623 bp in length containing a open reading frame of 1338 bp, and identical to the β -tubulin gene (Figure 1). This is the first CFTUB2 cDNA clone isolated from cotton. Comparing the nucleotide and predicted polypeptide sequences of the cotton CFTUB2 with the data banks, it was found that the CFTUB2 cDNA shared 96%-98% homology at the amino acid level and over 78% homology at the nucleotide level with the known β-tubulin cDNAs and genes from other plants (such as Arabidopsis, tobacco, rice, soybean, maize, potato, carrot, etc.) [Liaud et al., 1992; Snustad et al., 1992; Villemur et al, 1994; Tonoike, et al., 1994; Taylor et al., 1994; Kang et al., 1994; Okamura et al., 1997; Chu et al., 1998; Okamura et al., 19991.

Total RNAs from different tissues of cotton were used to reverse-transcribe first-strand cDNAs which were used as templates in differential display PCR reactions. Differential display analysis was carried out by using a differential display kit (Clontech). First-strand cDNA was synthesized with 2 pg total RNA as starting materials of reverse transcription and oligo(dT) as primers at 42°C for 1 hour. Differential display PCR reactions were carried out with a initial cycle consisting of 94°C for 5 minutes, 40°C for 5 minutes and 68°C for 5 minutes, followed by two cycles consisting of 94°C for 2 minutes and 40°C for 5 minutes and 68°C for 5 minutes, and then 25 cycles consisting of 94°C for 1 minute and 60°C for 1 minute and 68°C for 2 minutes, and a final extension at 68°C for 7 minutes. Target differential display bands were excised and reamplified for further analysis. Reproducible fiber-specific differential display products were targeted for further analysis. The cDNA in each target band was harvested and regenerated by PCR amplification. The isolated cDNA was subsequently subcloned into vectors and sequenced.

5

10

15

20

25

30

The Northern blot analysis showed that the transcripts of the CFTUB2 gene exhibited a highest accumulation in cotton young fibers of 8 DPA, and then there was a visible decrease in the accumulation of the gene products (mRNA) with further development of the fibers. Comparison of gene expression in different developmental stages of cotton ovules also showed that the gene transcripts accumulated more in 8 DPA ovules than in 4 and 14 DPA, and there was a gradual and visible decrease to an undetectable level in the accumulation of gene products with fiber development from 8 DPA to 28 DPA. This suggests that the gene is specifically expressed with a strict regulation at the transcriptional level during cotton fiber and ovule development, as seen with other cotton fiber-specific genes [Whittaker and

Triplett, 1999; Shin and Brown, 1999; Kawai et al., 1998; John, 1996a; Song and Allen, 1997; Ma et al, 1997; Rinehart et al., 1996; John and Crow, 1992].

Example 2: Isolation of the CFTUB2 promoter

5

10

15

20

25

30

Based on the screened CFTUB2 cDNA sequence, the CFTUB2 promoter was isolated from cotton Genome Walker libraries by Genome Walker PCR.

Total DNA was extracted and purified from leaves of cotton plants by using the following method. Liquid N_2 was added to 4 g of leaf tissues, and the leaves were homogenized thoroughly. 20 ml ice-cold extraction buffer (63 g/L glucose, 0.1 M Tris. HCl (pH 8.0), 5 mM EDTA, 20 g/L PVP-40, 1 g/L DIECA, 1 g/L ascorbic acid, 2ml/L, betamercaptoethanol) was added to the homogenized tissues in a 50 ml tube and centrifuged at 2500 rpm for 15 minutes. After removing the supernatant, 10 ml lysis buffer was added to each tube. The resuspended pellets were incubated at 65°C for 30 minutes. 10 ml chloroform was added to each tube, mixed with the samples and centrifuged at 3500 rpm for 10 minutes. The supernatant was transferred to a clean tube, and chloroform extraction was repeated one more time. The supernatant was transferred to a clean tube, and 0.6 volume isopropanol was added to each tube for DNA precipitation. After centrifuging at 3500 rpm for 30 minutes, the DNA was washed with 70% ethanol. The isolated genomic DNA was then dissolved in sterile water or TE (10 mM Tris.HCl, 1 mM EDTA) for use.

Cotton genomic DNA libraries were constructed from leaves of cotton plants. DNA was partially digested with BamH I, and the DNA fragments were cloned in the $\it BamH$ I site of the ZAP expression vector (Stratagene).

Genome Walker libraries were constructed by using Universal Genome Walker kit (Clontech). Genomic DNA from leaves

of cotton plants was digested with five restriction enzymes respectively, and then purified by phenol/chloroform and precipitated by ethanol. Digested DNA was ligated to Genome Walker adaptors. Two rounds of Genome Walker PCR reactions were carried out successively. 1 µl of each Genome Walker DNA library was used as templates in the primary PCR, and the primary PCR products were used as templates in secondary PCR. The PCR was started at 95°C for 1 minute, followed by 35 cycles consisting of 95°C for 15 seconds and 68°C for 4 minutes, and a final extension at 68°C for 6 minutes. Target PCR bands were cut out and purified by Geneclean kit (Bio 101).

5

10

15

20

25

3.0

Two fragments in the promoter region were isolated and cloned into pGEM-T vector, respectively. One fragment of the CFTUB2 promoter was 1433 bp in length Figure 2, and another was 984 bp long. Figure 2. A Hind III site and a BamH I site were created at the 5'-end and 3'-end of the 0.9 kb CFTUB2 promoter fragment of cotton respectively by PCR method. The Hind III/BamH I fragment was initially subcloned into pGEM-T vector (Promega). Plasmid DNA containing the CFTUB2 promoter fragment was digested with Hind III and BamH I, and the digested fragment was isolated by agarose gel electrophoresis. A chimeric CFTUB2 promoter/GUS construct was generated by insertion of the fragment, replacing CaMV 35S promoter, into the Hind III/BamH I sites of pBI121 vector.

The 1.3 kb of BamH I/BamH I CFTUB2 promoter fragment was initially subcloned into the pGEM-T vector (Promega). Plasmid DNA containing the CFTUB2 promoter fragment was digested with BamH I, and the digested fragment was isolated by agarose gel electrophoresis. A chimeric CFTUB2 promoter/GUS construct was generated by insertion of the fragment into the BamH I site of pBI101 vector.

Example 3: Functional analysis of the CFTUB2 promoter

5

10

1.5

20

25

30

In order to characterize the function of CFTUB2 promoter in fiber-specific expression of the CFTUB2 gene, a 1.3 kb of fragment and a 0.9 kb of fragment of the CFTUB2 promoter were fused with gus coding sequence in the gene expression vector pBI101 or pBI121 (deleting CaMV35S promoter), respectively (Figure 3). The constructs of CFTUB2 promoter/GUS fusion gene were used to transform tobacco and cotton by Agrobacterium-mediated gene transfer, using the pBI121 vector containing CaMV35S promoter/GUS fusion as a positive control. The CaMV35S promoter is active in all the tissues of cotton and other plants and is a constitutive promoter [Odell et al., 1985; Ow et al., 1987; McCabe and Martinell, 1993]. A binary vector containing either a CFTUB2 promoter/GUS fusion gene or the CaMV35S promoter/GUS control control was transferred into Agrobacterium tumefaciens strain LBA 4404. Cotton explants for transformation were obtained from cotton seedlings grown as in Example 1. Tobacco explant material was obtained from tobacco seedlings. Tobacco seeds were surface-sterilized with 70% ethanol for 30-60 seconds and 0.1% HgCl2 for 15 minutes, followed by washing with sterile water. The seeds germinated on ½ MS medium on light at 28°C in culture room, and leaves cut from sterile seedlings for use as explants for transformation. Cotton cotyledon and hypocotyl explants and tobacco leaf explants were transformed by the Agrobacterium with the vectors, and transformed plants were transplanted to soil in greenhouse for growing to maturity. -

Tobacco leaves were cut into about 2x2 cm pieces, and immersed in <u>Agrobacterium</u> suspension for 5 minutes. The infected tobacco explants were cultivated on MS medium with 1 mg/L 6-BA for 48 hours at 28°C, and then transferred onto selection MS medium containing 100 mg/L kanamycin and 1 mg/L 6-

BA for 20-30 days for selecting transformed shoots (kanamycin-resistant shoots). The transformed shoots were cut from the calli and rooted on MS medium with 50-100~mg/L kanamycin. The transformed tobacco plants were transplanted to soil in greenhouse for growing to maturity.

5

10

15

20

25

3.0

The cotyledon and hypocotyl were used as explants for cotton transformation. Cotton seeds were surface-sterilized with 70% ethanol for 30 seconds and 10% H₂O₂ for 60 minutes, followed by washing with steril water. These seeds were incubated in the sterile water at 28°C. After over night, the seeds sprouted. The embryos were taken out and put on the IM medium (1/2 (MS (macronutriants, micronutrients, EDTA-Fe) + VB1 10mg/L + VB6 1mg/L + VPP 1mg/L + Myo-Insitol 100mg/L} + phytogel 2q/L pH = 6.4) at $28^{\circ}C$ for 7 days. The cotyledon and hypocotyl of cotton were used as explants for transformation. After cutting into 5 mm2 (mm) piece, the explants were soaked in the Agrobacterium tumefaciens strain LBA 4404 suspension (OD₅₀₀ = 0.2 - 0.4) for 15 minutes. Then the explants were put on CM medium (MS (macronutriants, micronutrients, EDTA-Fe) + VB1 10mg/L + VB6 1mg/L + VPP 1mg/L + Myo-Instiol 100mg/L + 2.4-D 0.1mg/L + KT 0.1mg/L + Glucose 30g/L + MgCl₂ 0.7mg/L + phytogel 2q/L pH = 6.4) at 24 °C for 2 days. After washing with liquid MS medium, the explants were put on the SM medium (MS (macronutriants, micronutrients, EDTA-Fe) + VB1 10mg/L + VB6 $1mq/L = VPP \ 1mq/L + Myo-Insitol \ 100mq/L + 2.4-D \ 0.1mq/L + KT$ $0.1 \text{mg/L} + \text{Glucose } 30 \text{g/L} + \text{MgCl}_2 \cdot 0.7 \text{mg/L} + \text{phytogel } 2 \text{g/L} +$ Kanamycin 50mg/L + Cefutoxime 200mg/L pH = 6.4) on light at 28°C in culture room for selecting and the subculture was per month. After 2-3 months subculturing on SM, the calli were induced from explants. The calli were transferred on DM medium (MS (macronutriants, micronutrients, EDTA-Fe) + VB1 10mg/L + VB6 1mg/L + VPP 1mg/L + Myo-Insitol 100mg/L + KNO₃ 19g/L + MgCl₂

0.7mg/L + Glucose 30g/L + phytogel 3g/L pH = 6.4) and subcultured per month. After about 5 months, the somatic embryos begin to form. Continuing to culture the young embryos on DM medium until they develop into maturity. The mature embryos were transferred on GM medium (1/2 (MS (macronutriants, micronutrients, EDTA-Fe) + VBl 10mg/L + VB6 1mg/L + VPP 1mg/L + Myo-Insitol 100mg/L) + NAA 0.01mg/L + Glucose 30g/L + phytogel 3.5g/L pH = 6.4) in the box for developing into plantlets. And then the plantlets were transplanted in the soil for the plant growing and collecting the transgenic seeds.

Transgenic tobacco and cotton plants possessing the

5

10

15

20

25

3.0

chimeric CFTUB2 promoter/GUS gene (or 35S:GUS gene), and non-transformed plants as negative controls, were analyzed by DNA Southern blot hybridization and by GUS histochemical assay. Total genomic DNA from cotton and tobacco leaves were digested with restriction enzymes, separated on agarose gels, and transferred onto Hybond-N nylon membranes by capillary blotting. DNA Southern blots were hybridized in ExpressHyb solution (Clontech) at 68°C with 32P-DNA probes prepared by random labeling (Promega Prime-a-Gene Labeling System). After hybridization, the blots were washed at 68°C in $0.1 \times \text{SSC}$, 0.5%SDS for 30-60 minutes. The 32P-labeled nylon membranes were exposed to X-ray film at - 70°C for autoradiography. The results of Southern blot analysis demonstrated that CFTUB2 promoter/GUS gene was integrated into tobacco and cotton genomes. Total of 325 transformed cotton plants, which belong to 31 transformed lines, were obtained and transplanted in soil to grow to maturation

Histochemical assays for GUS activity in transgenic tobacco and cotton plants were conducted according to the protocol described previously by Jefferson et al. (1987) with some modifications. Fresh tissues from the plants were incubated

in X-gluc (5-bromo-4-chloro-3-indolvlglucuronide) solution consisting of 0.1 M sodium phosphate (pH 7.0), 10 mM ethylene diaminetetraacetic acid (EDTA), 0.5 mM potassium ferrocyanide and 0.5 mM potassium ferricyanide, and 0.1 % X-gluc (Clontech chemical) overnight. The stained plant materials were then cleared and fixed by rinsing with 100% and 70% ethanol successively, and the samples were examined and photographed directly or under a microscope. Consistent with the results from Northern blot analysis, the GUS gene driven by CFTUB2 promoter specifically expressed in the young fibers, but not in other tissues. in all the 31 transgenic cotton plants studied, while the GUS activity was detected in all the tissues of positive control cotton plants (35S:GUS). A total of 36 transformed cotton plants were obtained and transplanted in soil to grow to maturity, all of which had detectable GUS activity only in the young fibers, not in the flower organs such as anthers, petals and sepals, or in leaves and roots. Similarly, it was found that under the CFTUB2 promoter, GUS gene activity was only detected in the seeds in all of the 15 transgenic tobacco plants studied, suggesting the CFTUB2 promoter activity was also tissue-specific in tobacco. This result, together with the above Northern blot analysis, indicates that the CFTUB2 promoter controls gene specific expression at the transcriptional level in cotton fibers.

5

10

15

2.0

References

10

2.0

25

- An YQ, Huang S, McDowell JM, 1996. Conserved expression of the Arabidopsis ACT1 and ACT3 actin subclass in organ primordia and mature pollen. Plant Cell, 8(1):15-30.
- 5 Arthur JC, 1990. In Polymers: Fibers and Textile, A Compendium, ed. Kroschwitz, JI. (Wiley, New York), pp. 118-141.
 - Baird WV and Meagher RB, 1987. A complex gene superfamily encodes actin in petunia. EMBO J., 6(11):3223-31.
 - Basra AS and Malik CP, 1984. Development of the cotton fiber.

 Int. Rev. Cytol. 89:65-113.
 - Chu B, Wilson TJ, McCune-Zierath C, Snustad DP, Carter JV.,
 1998. Two beta-tubulin genes, TUB1 and TUB8, of
 Arabidopsis exhibit largely nonoverlapping patterns of
 expression. Plant Mol. Biol., 37(5):785-90.
- Cox GM, Rude TH, Dykstra CC, 1995. The actin gene from

 Cryptococcus neoformans: structure and phylogenetic

 analysis. J. Med. Vet. Mycol... 33(4):261-6.
 - Cyr RJ and Palevitz BA, 1995. Organization of cortical microtubules in plant cells. Curr. Opin. Cell Biol., 7:65-71.
 - Fisher DD and Cyr RJ, 1998. Extending the microtubule/ microfibril paradigm. Plant Physiol., 116:1043-51.
 - Giddings TH and Staehelin LA, 1991. Microtubule-mediated control of microfibril deposition: a re-examination of the hypothesis. In CW Lloyd, ed., The Cytoskeletal Basis of Plant Growth and Form. Academic Press, London, pp. 85-99.
 - Hasenfratz MP, Tsou CL, Wilkins TA, 1995. Expression of two related vacuolar H(+)-ATPase 16-kilodalton proteolipid genes is differentially regulated in a tissue-specific manner. Plant Physiol., 108(4): 1395-404.
 - Jefferson RA, 1987. Assaying chimeric genes in plants: the GUS
 gene fusion system. Plant Mol. Biol. Rep., 5:387-405.

- Jefferson RA, Kavanagh TA, Bevan MW, 1987. GUS fusion:
 p-glucuronidase as a sensitive and versatile gene fusion
 marker in higher plants. EMBO J., 6:3901
- John ME and Crow LJ, 1992. Gene expression in cotton fiber: cloning of the mRNAs. Proc. Natl. Acad. Sci. USA, 89(13):5769-73.

10

- John ME and Peterson MW, 1994. Cotton pollen-specific polygalacturonase mRNA: tissue and temporal specificity of its promoter in transgenic tobacco. Plant Mol. Biol., 26(6):1989-93.
- John ME, 1996a. Structural characterization of genes corresponding to cotton fiber mRNA, E6: reduced E6 protein in transgenic plants by antisense gene. Plant Mol. Biol., 30(2):297-306.
- John, ME, 1996b. Genetically engineering cotton plants for altered fiber, U.S. Patent No. 5,495,070.
 - John, ME, and Keller, G. 1996. Metabolic pathway engineering in cotton: Biosynthesis of polyhydroxybutratein fiber cells. Proc. Natl. Acad. Sci. USA. 93:12768-12773.
- John, ME, 1997a. Transgenic cotton plants producing heterologous polyhydroxy(e) butyrate bioplastic, U.S. Patent No. 5,602,321.
 - John, ME, 1997b. Genetically engineering cotton plants for altered fiber, U.S. Patent No. 5,620,882 (1997).
- 25 Kang MS, Choi YJ, Kim MC, Lim CO, Hwang I, Cho MJ., 1994.
 Isolation and characterization of two beta-tubulin cDNA clones from rice. Plant Mol. Biol., 26(6): 1975-9.
 - Kawai M, Aotsuka S, Uchimiya H, 1998. Isolation of a cotton CAP gene: a homologue of adenylyl cyclase-associated protein highly expressed during fiber elongation. Plant Cell Physiol., 39(12):1380-3.

- Liaud MF, Brinkmann H, Cerff R., 1992. The beta-tubulin gene family of pea: primary structures, genomic organization and intron-dependent evolution of genes. Plant Mol. Biol., 18(4):639-51.
- 5 Loguerico LL, Zhang JQ, Wilkins TA, 1999. Differential regulation of six novel MYB-domain genes def two distinct expression patterns in allotetraploid cotton. Mol. Gen. Genet., 261(4/5):660-71.
 - Ma DP, Liu HC, Tan H, Creech RG, Jenkins JN, Chang YF, 1997.

 Cloning and characterization of a cotton lipid transfer protein gene specifically expressed in fiber cells.

 Biochim. Biophys. Acta. 1344(2): 111-4.

15

20

- McCabe DE and Martinell BJ, 1993. Transformation of elite cotton cultivars via particle bombardment of meristems.

 Biotechnology, 11:596-8.
- McElroy D, Rothenberg M, Reece KS, Wu R, 1990. Characterization of the rice actin gene family. Plant Mol. Biol., 15(2):257-68.
- Nairn CJ, Winesett L, Ferl RJ, 1988. Nucleotide sequence of an actin gene from Arabidopsis thaliana. Gene, 65(2):247-57.
- Odell JT, Nagy F, Chua N-H, 1985. Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. Nature, 313:810-2.
- Okamura S, Naito K, Sonehara S, Ohkawa H, Kuramori S, Tatsuta M, Minamizono M, Kataoka T., 1997. Characterization of the carrot beta-tubulin gene coding a divergent isotype, beta-2. Cell Struct. Funct., 22(2):291-8.
- Okamura S, Okahara K, Iida T, Ozaki M, Asano S, Morita M,
 Imanaka T, 1999. Isotype-specific changes in the amount
 of beta-tubulin RNA in synchronized tobacco BY2 cells.
 Cell Struct. Funct., 24(3): 117-22.

- Ow DW, Jacobs JD, Howell SH, 1987. Functional regions of the cauliflower mosaic virus 35S RNA promoter determined by use of the firefly luciferase gene as a reporter of promoter activity. Proc. Natl. Acad. Sci. USA, 84:4870-4.
- 5 Rinehart JA, Peterson MW, John ME, 1996. Tissue-specific and developmental regulation of cotton gene FbL2A. Demonstration of promoter activity in transgenic plants. Plant Physiol., 112(3):1331-41.
 - Ryser U, 1985. Cell wall biosynthesis in differentiating cotton fibers. Eur. J. Cell Biol., 39:236-56.
 - Shah DM, Highrower RC, Meagher RB, 1983. Genes encoding actin in higher plants: intron positions are highly conserved but the coding sequences are not. J. Mol. Appl. Genet., 2(1): 111-26
- 15 Shin H and Brown RM jr, 1999. GTPase activity and biochemical characterization of a recombinant cotton fiber annexin. Plant Physiol., 119(3):925-34.

20

25

- Song P and Allen RD, 1997. Identification of a cotton fiberspecific acyl carrier protein cDNA by differential display. Biochim. Biophys. Acta, 1351(1):305-12.
- Snustad DP, Haas NA, Kopczak SD, Silflow CD., 1992. The small genome of Arabidopsis contains at least nine expressed beta-tubulin genes. Plant Cell, 4(5):549-56.
- Stranathan M, Hastings C, Trinh H, 1989. Molecular evolution of two actin genes from carrot. Plant Mol. Biol., 13(4):375-83.
 - Taylor MA, Wright F, Davies HV., 1994. Characterization of the cDNA clones of two beta-tubulin genes and their expression in the potato plant (Solanum tuberosum L.). Plant Mol. Biol., 26(3):1013-18.

- Tonoike H, Han IS, Jongewaard I, Doyle M, Guiltinan M, Fosket

 DE, 1994. Hypocotyl expression and light downregulation

 of the soybean tubulin gene, tubB1. Plant J.,

 5(3):343-51.
- 5 Umbeck, Paul, 1992. Genetic engineering of cotton plants and lines, U.S. Patent No. 5,159,135.

- Villemur R, Haas NA, Joyce CM, Snustad DP, Silflow CD, 1994.

 Characterization of four new beta-tubulin genes and their expression during male flower development in maize (Zea mays L.). Plant Mol. Biol., 24(2):295-315.
- Whittaker DJ and Triplett BA, 1999. Gene-specific changes in alpha-tubulin transcript accumulation in developing cotton fibers. Plant Physiol., 121(1):181-8.

We claim:

- 1 1. A promoter that is cotton fiber-specific, comprising the promoter of the cotton β -tubilin gene CFTUB2.
- 1 2. A promoter that is cotton fiber-specific, comprising a 1433 kb fragment of the promoter of the cotton β -tubilin gene CFTUB2 having the sequence of SEQ ID NO: 2.
- 1 3. A promoter that is cotton fiber-specific, comprising a 2 984 kb fragment of the promoter of the cotton β -tubilin 3 gene CFTUB2 having the sequence of nucleotides 449-1433 of SEQ ID NO: 2.

ABSTRACT

The present invention relates to a cotton β -tubulin gene CFTUB2, and active fragments thereof. These promoters show strong fiber-specific activity.

5 2577-138.AP1

10 Review 10089543 - 080402

. 2577-138 MTB:clw

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of Lin CAI et al. Serial No. 10/089,543 Examiner: Filed: April 1, 2002) Group Art Unit:

For: ISOLATION AND CHARACTERIZATION OF A FIBER-SPECIFIC B-TUBULIN

PROMOTER FROM COTTON

SUBMISSION OF SEQUENCE LISTING

Assistant Commissioner for Patents Washington, D.C. 20231

Dear Sir:

Attached is a paper copy and a computer readable copy of a Sequence Listing in the above-referenced application. Applicants certify that the sequence listing information recorded in computer readable form is identical to the paper copy of the sequence listing and contains no new matter.

Respectfully submitted,

Mark I. Bowditch

Attorney for Applicants Registration No. 40,315 ROTHWELL, FIGG, ERNST & MANBECK, p.c. Suite 800, 1425 K Street, N.W.

Washington, D.C. 20005 Telephone: (202) 783-6040

2577-138.sub

10089543.080402 \$9.589.543

10 Receipt 0 5 AUG 2002

SEQUENCE LISTING

<110> Cai, Lin Li, Xuebao Cheng, Ninghui Liu, Jian-Wei

<120> Isolation and Characterization of a Fiber-Specific b-Tubulin Promoter from Cotton

<130> 2577-138

<140> 10/089543

<141> 2002-04-01

<150> PCT/SG00/00111

<151> 2000-08-01

<160> 2

<170> PatentIn version 3.1

<210> 1

<211> 1623 <212> DNA

<213> Arabidopsis sp.

<400> ggcacgagta tatttttcct ctccaatttt ccqtcacttt cccqaqaaaa tqaqaqaaat 60 120 cottcacate caaggtggce aatgeggcaa teagatagga gecaagttet gggaagtegt atgtgccgaa catggcatcg attcaacggg tcgatatggt ggtgactcgg agctccaqct 180 tgagcgaatc aatgtttact acaacgaagc cagttgtggc cgttttgttc cccgcgcagt 240 tttaatggat ctggaacccg gaaccatgga tagcgtaaga tccggtcctt acggacaaat 300 tttccqaccc gataacttcg ttttcggaca gtccggtgcg ggaaacaatt gggctaaggg 360 acattacact gaaggagcgg agcttatcga ttccgttctc gacgtggtta gaaaggaagc 420 cqaaaattqc gattgcttgc aagggtttca ggtatgccat tctttgggaa gaagaacggg 480 ttccqqaatq ggaacgttgt tgatatcgaa gatacgagag gagtatccgg accgaatgat 540 gettaegttt teggtgttte cateteceaa ggtttetgat aetgttgttg aacettaeaa 600 cqcqacactc tcaqttcatc aqcttqtqqa aaatqctgat qaqtqtatqq ttcttqataa 660 cgaagetete tacgatatet gttteegtae eetcaagete actaetecaa gttttggaga 720

780 totcaaccat ctaatttotg ccaccatgag tggtgtaaca tgctgccttc gcttccctgg 840 tttcttcatg gtgggatttg cgcctctcac ctcacgcgqt tcccaacaqt acaqaqccct 900 960 cactgtccct gaacttacac agcaaatgtg ggatgccaag aacatgatgt gtgcagctga tectequeae qqtequatace teacaqeate ageggtette egtggquaqu tquqeaequa 1020 1080 agaggttgat gagcagatga tcaatgtgca aaacaagaac tcatcttact ttgttgaatg qatcccqaac aatqtqaaqt ccactgtttg tgacatccct ccaatcggct taaagatggc 1140 atccacattt atcgggaact ctacttcaat ccaagagatg ttcaggaggg tgagtgaaca 1200 attcactgcc atgttccgta ggaaagcttt cttgcattgg tatactggag aagggatgga 1260 tgagatggag ttcacagaag cggagagtaa catgaatgac ttggtttctg agtaccaaca 1320 ataccaqqat qcaactqcaq atgatgaaga gtacgaggaa gaggaggaat acgaggcaga 1380 qqcttaaaat ctaatqqaat aatttggatg tttttcgttg tgttttggat tqqqcttgtg 1440 qaqtqtqttq atqcaatttc tcactqcctq tttttqgtct ttggatcact qtattgttga 1500 1560 ttgtgtggac tttagttttg tcctcacagc ttacggagta tatgttgttg tattgcttgt 1620 1623 aaa

<400> 2

actataggge acgegggte gacggecegg getggtgett gatatetatg attteagat 60
ttgcataaga ettetateta teagaagaeg cetgeagagg ateceaaatt agtetaaaat 120
tatetteagt eteggaaace aacteaggae ecaaaaceeg tegeteacee aacteagtet 180
aatataacag agtatgacae ttatgaceat atagageete gtaaggtgee atetagatge 240
cagattggaa actgttattg taggegaact caactaacegg taaaaaatee teteaactae 300
cttagtaata aateacatag etecaaateg tateetetag tatatgaate acetteetaa 360
attgaceate ggtetgagga tggaatgeag aceggtgeea eegatttaet aatggtaeet 420

<210> 2 <211> 1

<211> 1433

<212> DNA

<213> Arabidopsis sp.

304030.E420800L

ataaaaaatt	attattttt	aaaaaattga	tgtgaccagt	ggttggagag	agaggtctac	480
cgattggtca	agtggcacca	atttttatt	ttacctcctg	cctagattcg	taaatactat	540
tgcatttatc	tcatttcatt	atttatttaa	ttattttata	ttatttggat	aaaaattcta	600
atactttact	tttttttaaa	aagaatttat	ttaattattt	tatattattt	agataaaaat	660
tctaatactt	tactttttt	ttaaaaagaa	tttcaattgc	gttttttctt	aatttagttt	720
taattctata	ctaattataa	aaattctgat	cggattagtg	tggtgtcaaa	gtcaagtcac	780
atgaattttg	ttggagaaaa	aataaaaatt	aaacacattt	ttcgattaat	ttattatata	840
tataataata	taaacacatt	tttatttaat	gttgtcaata	atattttta	attaaaattt	900
cagcacaaca	attacactct	catcattaaa	tttaatctta	ttaccataat	taaaattgtg	960
aggacaatta	ttttttaatc	tcaccctcca	ttaatgcata	ttattaattt	ttgttcgata	1020
cttcttattt	cactcctaac	attaatcatt	aacccaattt	tgaactgtta	taatttetta	1080
acttattcac	tattgtggct	ctgggtccat	ctggaaaggc	caccgtccag	gctgtccaac	1140
cacactttgc	cacgtcatca	attccagtaa	ctacattgtt	acagttacta	agcaaatccc	1200
aatttcaaaa	attcaatttc	ccaggaaaac	gaaacgtccg	ttactaaccg	acctaaaacc	1260
cagctcaacc	tgccgtcaat	taacggaaat	cttttaactc	ctctatataa	cccaaaacca	1320
ctctcatcac	catttcccca	taaaaagaat	ttccggaatt	cttattcctt	ttatattttt	1380
cctctccaat	ttcccgtcac	tttccggaga	aaatgagaga	aatccttcac	atc	1433

Inventor: Lin CAI et al.

ISOLATION AND CHARACTERIZATION OF A FIBER-SPECIFIC
β-TUBULIN PROMOTER FROM COTTON

Docket No. 2577-138

GGCACGAGTATATTTTTCCTCTCCAATTTTCCGTCACTTTCCCGAGAAAATGAGAGAAATC CTTCACATCCAAGGTGGCCAATGCGGCAATCAGATAGGAGCCAAGTTCTGGGAAGTCGTA TGTGCCGAACATGGCATCGATTCAACGGGTCGATATGGTGGTGACTCGGAGCTCCAGCTT GAGCGAATCAATGTTTACTACAACGAAGCCAGTTGTGGCCGTTTTGTTCCCCGCGCAGTTT TAATGGATCTGGAACCCGGAACCATGGATAGCGTAAGATCCGGTCCTTACGGACAAATTT TCCGACCCGATAACTTCGTTTTCGGACAGTCCGGTGCGGGAAACAATTGGGCTAAGGGAC ATTACACTGAAGGAGCGGAGCTTATCGATTCCGTTCTCGACGTGGTTAGAAAGGAAGCCG AAAATTGCGATTGCTTGCAAGGGTTTCAGGTATGCCATTCTTTGGGAAGAAGAACGGGTT CCGGAATGGGAACGTTGTTGATATCGAAGATACGAGAGGAGTATCCGGACCGAATGATGC TTACGTTTTCGGTGTTTCCATCTCCCAAGGTTTCTGATACTGTTGTTGAACCTTACAACGCG ACACTCTCAGTTCATCAGCTTGTGGAAAATGCTGATGAGTGTATGGTTCTTGATAACGAAG CTCTCTACGATATCTGTTTCCGTACCCTCAAGCTCACTACTCCAAGTTTTGGAGATCTCAAC CATCTAATTTCTGCCACCATGAGTGGTGTAACATGCTGCCTTCGCTTCCCTGGTCAGCTTA ACTCAGATCTCCGCAAACTTGCTGTAAACCTTATTCCATTCCCTCGACTACATTTCTTCATG GTGGGATTTGCGCCTCTCACCTCACGCGGTTCCCAACAGTACAGAGCCCTCACTGTCCCTG AACTTACACAGCAAATGTGGGATGCCAAGAACATGATGTGTGCAGCTGATCCTCGACACG GTCGATACCTCACAGCATCAGCGGTCTTCCGTGGGAAGATGAGCACGAAAGAGGTTGATG AGCAGATGATCAATGTGCAAAACAAGAACTCATCTTACTTTGTTGAATGGATCCCGAACA ATGTGAAGTCCACTGTTTGTGACATCCCTCCAATCGGCTTAAAGATGGCATCCACATTTAT CACAGAAGCGGAGAGTAACATGAATGACTTGGTTTCTGAGTACCAACAATACCAGGATGC AACTGCAGATGATGAAGAGTACGAGGAAGAGGGGGAGTACGAGGCAGAGGCTTAAAATC TAATGGAATAATTTGGATGTTTTTCGTTGTTTTTGGATTGGGCTTGTGGAGTGTTTTGATG CAATTTCTCACTGCCTGTTTTTGGTCTTTGGATCACTGTATTGTTGATTGTTGTCGACTTTAG TTTTGTCCTCACAGCTTACGGAGTATATGTTGTTGTATTGCTTGTTGATTCATCTTATAAGT

Inventor: Lin CAI et al.
ISOLATION AND CHARACTERIZATION OF A FIBER-SPECIFIC
\$\text{\text{\$\text{\$\text{\$PTUBULIN PROMOTER FROM COTTON}\$}}}
Docket No. 2577-138

ACTATAGGGCACGCGTGGTCGACGGCCCGGGCTGGTGCTTGATATCTATGATTTTCAGATT TGCATAAGACTTCTATCTATCAGAAGACGCCTGCAGAGGATCCCAAATTAGTCTAAAATT ATCTTCAGTCTCGGAAACCAACTCAGGACCCAAAACCCGTCGCTCACCCAACTCAGTCTA ATATAACAGAGTATGACACTTATGACCATATAGAGCCTCGTAAGGTGCCATCTAGATGCC AGATTGGAAACTGTTATTGTAGGCGAACTCAACTAACGGTAAAAAATCCTCTCAACTACC TTAGTAATAAATCACATAGCTCCAAATCGTATCCTCTAGTATATGAATCACCTTCTCAAAT TGACCATCGGTCTGAGGATGGAATGCAGACCGGTGCCACCGATTTACTAATGGTACCTAT ATTGGTCA AGTGGCACCAATTTTTTATTTTACCTCCTGCCTAGATTCGTAAATACTATTGCA ACTTTTTTTTAAAAAGAATTTATTTAATTATTTTATATTTTTAGATAAAAATTCTAATACT TTACTTTTTTTAAAAAGAATTTCAATTGCGTTTTTTCTTAATTTAGTTTTAATTCTATACT AATTATAAAAATTCTGATCGGATTAGTGTGGTGTCAAAGTCAAGTCACATGAATTTTGTTG CTCTCATCATTAAATTTAATCTTATTACCATAATTAAAATTGTGAGGACAATTATTTTTTAA TCTCACCCTCCATTAATGCATATTATTAATTTTTTGTTCGATACTTCTTATTTCACTCCTAACA TTAATCATTAACCCAATTTTGAACTGTTATAATTTCTTAACTTATTCACTATTGTGGCTCTG GGTCCATCTGGAAAGGCCACCGTCCAGGCTGTCCAACCACACTTTGCCACGTCATCAATTC CAGTAACTACATTGTTACAGTTACTAAGCAAATCCCAATTTCAAAAATTCAATTTCCCAGG AAAACGAAACGTCCGTTACTAACCGACCTAAAACCCAGCTCAACCTGCCGTCAATTAACG GAATTTCCGGAATTCTTATTCCTTTATATTTTTCCTCCCAATTTCCCGTCACTTTCCGGAG A A A A TGA GA GA A A TCCTTCACATC

Inventor: Lin CAI et al.
ISOLATION AND CHARACTERIZATION OF A FIBER-SPECIFIC
β-TUBULIN PROMOTER FROM COTTON
Docket No. 2577-138

CFTUB2 gene

cDNA (ORF: 1338 bp	

pBITUB vectors
0.9 kb

1.3 kb

CFTUB2 promoter gus gene NOS



JC13 Rec'd PCT/PTC 0 1 APR 2002

SEQUENCE LISTING

```
<110> Cai, L.
     Li, X. B.
      Chen, N. H.
     Liu, J. W.
<120> Isolation and Characterization of a Fiber-Specific
      Beta-Tubulin Promoter from Cotton
<130> 2577-138
<140>
<141>
<160> 2
<170> PatentIn Ver. 2.1
<210> 1
<211> 1623
<212> DNA
<213> Arabidopsis sp.
<400> 1
ggcacgagta tatttttcct ctccaatttt ccgtcacttt cccgagaaaa tgagagaaat 60
cettcacate caaggtggee aatgeggeaa teagatagga gecaagttet gggaagtegt 120
atgtgccgaa catggcateg attcaacggg tcgatatggt ggtgactcgg agctccaqct 180
tgagcgaatc aatgittact acaacgaagc cagttgtggc cgttttgttc cccgcgcagt 240
tttaatggat etggaaceeg gaaceatgga tagegtaaga teeggteett aeggacaaat 300
tttccgaccc gataacttcg ttttcggaca gtccggtgcg ggaaacaatt gggctaaggg 360
acattacact gaaggagegg agettatega tteegttete gaegtggtta gaaaggaage 420
cgaaaattgc gattgcttgc aagggtttca ggtatgccat tetttgggaa gaagaacggg 480
ttccggaatg ggaacgttgt tgatatcgaa gatacgagag gagtatccgg accgaatgat 540
gettacgttt teggtgttte cateteccaa ggtttetgat actgttgttg aacettacaa 600
cgcgacactc tcagttcatc agcttgtgga aaatgctgat gagtgtatgg ttcttgataa 660
cgaagetete tacgatatet gitteegtae ecteaagete actaetecaa gittiggaga 720
tetcaaccat ctaatttetg ccaccatgag tggtgtaaca tgctgcette gettecetgg 780
tcagcttaac tcagatetec gcaaacttgc tgtaaacctt attccattcc ctcgactaca 840
tttcttcatg gtgggatttg cgcctctcac ctcacgcggt tcccaacagt acagagccct 900
cactgtccct gaacttacac agcaaatgtg ggatgccaag aacatgatgt gtgcagctga 960
tcctcgacac ggtcgatacc tcacagcatc agcggtcttc cgtgggaaga tgagcacgaa 1020
agaggttgat gagcagatga tcaatgtgca aaacaagaac tcatcttact ttgttgaatg 1080
gatcccgaac aatgtgaagt ccactgtttg tgacatccct ccaatcggct taaagatggc 1140
atccacattt atcgggaact ctacttcaat ccaagagatg ttcaggaggg tgagtgaaca 1200
attcactgcc atgitccgta ggaaagcttt cttgcattgg tatactggag aagggatgga 1260
tgagatggag ttcacagaag cggagagtaa catgaatgac ttggtttctg agtaccaaca 1320
ataccaggat gcaactgcag atgatgaaga gtacgaggaa gaggaggaat acgaggcaga 1380
ggcttaaaat ctaatggaat aatttggatg tttttcgttg tgttttggat tgggcttgtg 1440
gagtgtgttg atgcaattte teactgeetg tttttggtet ttggateact gtattgttga 1500
ttgtgtcgac tttagttttg tectcacage ttacggagta tatgttgttg tattgettgt 1560
aaa
```



<210> 2 <211> 1433 <212> DNA <213> Arabidopsis sp.

<400> 2

actatagggc acqcgtggtc qacqgcccgg gctggtgctt gatatctatg attttcagat 60 ttqcataaqa cttctatcta tcagaaqacg cctgcagagg atcccaaatt agtctaaaat 120 tatetteagt eteggaaace aacteaggae ceaaaaceeg tegeteacee aacteagtet 180 aatataacag agtatgacac ttatgaccat atagagcctc gtaaggtgcc atctagatgc 240 cagattggaa actgttattg taggcgaact caactaacgg taaaaaatcc tctcaactac 300 cttagtaata aatcacatag ctccaaatcg tatcctctag tatatgaatc accttctcaa 360 attgaccate ggtetgagga tggaatgeag aceggtgeea eegatttaet aatggtacet 420 ataaaaaatt attattttt aaaaaattga tgtgaccagt ggttggagag agaggtctac 480 cqattqqtca aqtgqcacca attttttatt ttacctcctg cctaqattcq taaatactat 540 tqcatttatc tcatttcatt atttatttaa ttatttata ttatttgqat aaaaattcta 600 atactttact tttttttaaa aagaatttat ttaattattt tatattattt agataaaaat 660 totaataott taottttttt ttaaaaaqaa tttoaattgo qttttttott aatttaqttt 720 taattotata otaattataa aaattotgat oggattagtg tggtgtcaaa gtcaagtcac 780 atgaattttg ttggagaaaa aataaaaatt aaacacattt ttcgattaat ttattatata 840 tataataata taaacacatt tttatttaat gttgtcaata atatttttta attaaaattt 900 caqcacaaca attacactot catcattaaa tttaatotta ttaccataat taaaattgtg 960 aggacaatta ttttttaatc tcaccctcca ttaatgcata ttattaattt ttgttcgata 1020 cttcttattt cactcctaac attaatcatt aacccaattt tgaactgtta taatttctta 1080 acttattcac tattqtqqct ctggqtccat ctggaaaggc caccgtccag gctgtccaac 1140 cacactttgc cacgtcatca attccagtaa ctacattgtt acagttacta agcaaatccc 1200 aatttcaaaa attcaatttc ccaggaaaac gaaacgtccg ttactaaccg acctaaaacc 1260 cageteaace tgccgteaat taacggaaat ettttaacte etetatataa eecaaaacea 1320 ctctcatcac catttcccca taaaaagaat ttccggaatt cttattcctt ttatattttt 1380 cctctccaat ttcccgtcac tttccggaga aaatgagaga aatccttcac atc

			Attorney Docket No.	2577-138
DECLARATION AND POWER OF		First Named Inventor	Lin CAI	
ATTORNEY FOR UTILITY OR DESIGN		COMPLETE IF KNOWN		
PATENT APPLICATION (37 CFR 1.63)		Application Number	PCT/SG00/00111	
Submitted Submitted	Declaration	Filing Date	August 1, 2000	
		Submitted after Initial Filing	Group Art Unit	Unassigned
	Filing		Examiner Name	Unassigned

As a below named inventor. I hereby declare that:

My residence, mailing address, and citizenship are as stated below next to name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: ISOLATION AND CHARACTERIZATION OF A FIBER-SPECIFIC.FTUBULIN PROMOTER FROM COTTON, the specification of which is was filed on August 1, 2000 as PCT International Application Number PCT/SG00/00111 and was amended on .

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application	Country	Foreign Filing Date	Priority Not	Certified Co	py Attached?
Numbers		(MM/DD/YYYY)	Claimed	YES	NO

hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.					
Application Number(s)	Filing Date (MM/DD/YYYY)				

I or we hereby appoint the registered practitioner(s) associated with Customer No. 6449 to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith. Direct all correspondence to Customer Number 6449.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR: [] A petition has been filed for this unsigned inventor					
Given Name <u>Lin</u> (first and middle [if any])		Family Name CAI or Surname			
Inventor's Signature	in	Date 12 - 04 - 2002			
Residence: City	State	Country Singapore	Citizenship China		
Mailing Address Blk 715, #07-79, Clementi	West St. 2, Singapore 1	20715			
NAME OF SECOND INVENTOR:	[] A petition ha	as been filed for this uns	igned inventor		
Given Name Xuebao (first and middle [if any])		Family Name_LI or Surname			
Inventor's Signature a Xueba	Date 12-04-2002				
Residence: City	State	Country Singapore			
Mailing Address 50 West Coast Crescent, #0	4-04, Westbay Condom	inium, Singapore 12803	5		
NAME OF THIRD INVENTOR:	as been filed for this unsigned inventor				
Given Name Ninghui (first and middle [if any])		Family Name CHENG or Surname			
Inventor's Signature		Date			
Residence: City Houston	State TX	Country U.S.A.	Citizenship China		
Mailing Address 7490 Brompton Road, Apt.	128, Houston, TX 7702	5			
City Houston	State TX	Zip 77025	Country U.S.A.		
NAME OF FOURTH INVENTOR:	[] A petition	has been filed for this ur	nsigned inventor		
Given Name Jian-Wei (first and middle [if any])		Family Name LIU or Surname			
Inventor's Signature		Date			
Residence: City	State	Country Singapore	Citizenship Singapore		
Mailing Address 8 Toh Yi Drive, #03-299, Singapore 590008					

NAME OF SOLE OR FIRST INVENTOR:	nas been filed for this unsigned inventor				
Given Name Lin (first and middle [if any])	Family Name CAI or Surname				
Inventor's Signature	0	Date			
Residence: City	State	Country Singapore	Citizenship China		
Mailing Address Blk 715, #07-79, Clementi	West St. 2, Singapore 12	20715			
NAME OF SECOND INVENTOR:	[] A petition ha	as been filed for this uns	igned inventor		
Given Name Xuebao (first and middle [if any])		Family Name LI or Surname			
Inventor's Signature		Date			
Residence: City	State	Country Singapore	Citizenship China		
Mailing Address 50 West Coast Crescent, #04-04, Westbay Condominium, Singapore 128035					
NAME OF THIRD INVENTOR: [] A petition has been filed for this unsigned inventor			signed inventor		
Given Name Ninghui (first and middle [if any])		Family Name CHENG or Surname			
Inventor's Signature		Date 4/18/02			
Residence: City Houston	State TX	Country U.S.A.	Citizenship China		
Mailing Address 7490 Brompton Road, Apt.	128, Houston, TX 7702	5			
City Houston	State TX	Zip 77025	Country U.S.A.		
NAME OF FOURTH INVENTOR: [] A petition		has been filed for this unsigned inventor			
Given Name Jian-Wei (first and middle [if any])		Family Name LIU or Surname			
Inventor's Signature		Date			
Residence: City	State	Country Singapore	Citizenship Singapore		
Mailing Address 8 Toh Vi Drive #03-299 Singapore 590008					



65 77,34988

NAME OF SOLE OR FIRST INVENTOR: [] A petition has been filed for this unsigned inventor					
Given Name Lin (first and middle [if any])		Family Name CAI or Surname			
Inventor's Signature		Date			
Residence: City	State	Country Singapore	Citizenship China		
Mailing Address Blk 715, #07-79, Clementi	West St. 2, Singapore 1	20715			
NAME OF SECOND INVENTOR:	[] A petition h	as been filed for this uns	signed Inventor		
Given Name Xuebao (first and middle [if any])		Family Name LI or Surname			
Inventor's Signature		Date			
Residence: City	State	Country Singapore	Citizenship China		
Mailing Address 50 West Coast Crescent, #04-04, Westbay Condominium, Singapore 128035					
NAME OF THIRD INVENTOR:	as been filed for this unsigned inventor				
Given Name Ninghui (first and middle [if any])		Family Name CHENG or Surname	Family Name CHENG or Surname		
Inventor's Signature		Date			
Residence: City Houston	State TX	Country U.S.A.	Citizenship China		
Mailing Address 7490 Brompton Road, Apt.	128, Houston, TX 7702	25			
City Houston	State TX	Zip 77025	Country U.S.A.		
NAME OF FOURTH INVENTOR:	[] A petition	has been filed for this u	nsigned inventor		
Given Name Jian-Wei (first and middle [if any])		Family Name LIU or Sumame			
Inventor's Signature		Date April	(17,2002		
Residence: City	-State	Country Singapore	CitizenshIp Singapore		
Mailing Address 8 Toh Yi Drive. #03-299, Singapore 590008					

